

**LUSAKA**  
CELL HELPLINE:  
0975 835581  
0955 835581



**KITWE**  
CELL HELPLINE:  
0971 257674  
0966 465017

## **HYBRID POULTRY FARM (Z) LTD**

### **BROILER VACCINATION GUIDE**

<b>AGE (DAYS)</b>	<b>DISEASE</b>	<b>DATE DUE</b>	<b>DATE COMPLETED</b>	<b>METHOD OF ADMINISTRATION</b>
<b>10 DAYS</b>	<b>GUMBORO</b>			<b>DRINKING WATER</b>
<b>14 DAYS</b>	<b>NEWCASTLE DISEASE</b>			<b>DRINKING WATER</b>
<b>18 DAYS</b>	<b>GUMBORO</b>			<b>DRINKING WATER</b>
<b>21 DAYS</b>	<b>NEWCASTLE DISEASE</b>			<b>DRINKING WATER</b>

#### ***PRACTICAL TIPS FOR VACCINATION***

- 1. DAY 9- START THE BIRDS ON VITAMINS IN READINESS FOR VACCINATION ON THE 10TH DAY**
- 2. NEVER VACCINATE SICK BIRDS**
- 3. THE AMOUNT OF WATER TO BE USED SHOULD BE EQUAL TO THE QUANTITY DRUNK WITHIN A 2 HR PERIOD THE DAY BEFORE VACCINATION**
- 8. TO STIMULATE THIRST, WITHDRAW WATER FOR 1 TO 2 HRS BEFORE VACCINATION, DEPENDING ON THE SEASON OF THE YEAR**
- 4. ALWAYS FOLLOW THE INSTRUCTIONS OF THE VACCINE MANUFACTURER AND CHECK EXPIRY DATES.**
- 5. AVOID FREEZING THE VACCINE - STORE BETWEEN 2-8 oC**
- 6. AVOID STORING VACCINE IN DIRECT SUNLIGHT**
- 7. ADD 2 GRAMS SKIMMED MILK PER LTR OF WATER AND ALLOW 30 MINUTES BEFORE ADDING THE VACCINE TO THE WATER**
- 8. REMOVE ALUMINIUM SEAL FROM THE VIAL**
- 9. REMOVE THE RUBBER STOPPER UNDER THE PREPARED WATER AND HALF FILL VIAL WITH MILK WATER, SHAKE GENTLY TO DISOLVE, POUR BACK INTO THE MIXTURE AND DISTRIBUTE.**
- 10. AFTER 1-1<sup>1</sup>/<sub>2</sub> HRS, WITHDRAW WHAT IS LEFT. CLEAN THE DRINKERS, GIVE FRESH WATER LACED WITH VITAMINS (IMMUNOTRESS).**

[www.hybridpoultryfarm.com](http://www.hybridpoultryfarm.com)

Email: [sales@hybridpoultryfarm.com](mailto:sales@hybridpoultryfarm.com)

The above guide gives our customers a suggested vaccination program and Hybrid Poultry will not be held liable for any claims in regards to this guideline.

Hybrid Poultry Farm suggest customers consult with their independant veterinarian for recomended application days, vaccines and dosages.

