


COMMERCIAL LAYER VACCINATION GUIDE

HYBRID BROWN LAYER-2020

COMMERCIAL LAYER FLOCK HEALTH MONITORING GUIDE

LAYER FLOCK N:

					HYBRID POULTRY FARM (Z) LTD		Placement date		
					BOX 310097				
					CHAMBA VALLEY - LUSAKA - ZAMBIA				Flock Number:
									Layers House N.
				Index Date					
Days	Weeks	Date-Due	Date-Done	Type	Details	Vaccine	Expiry	Document	
1	0.1			V	Rispens CVI 988 HATCHERY LEVEL-MAREK'S SUBCUT			Done	
1	0.1	1-Jan-00		V	VH+H120: HATCHERY LEVEL-SPRAY NCD+IBV			Done	
10	1.3	10-Jan-00		V	GUMBORO MB: Drinking Water				
13	1.8	13-Jan-00		V	IB 4/91: Spray Vaccination Infectious Bronchitis				
18	2.4	1.8-Jan-00		V	GUMBORO MB: Drinking Water				
21	3	21-Jan-00		V	VH: Spray Vaccination NCD				
49	7	19-Feb-00		V	Ma5+Clone30: Spray Vaccination NCD+IBV				
56	8	25-Feb-00		V	CEO POX + Dil: Wing Web - full				
70	10	10-Mar-00		V	Nobilis IB + ND: I.M.0.5ml INAC				
77	11	17-Mar-00		V	IB 4/91: Spray Vaccination Infectious Bronchitis				
84	12	24-Mar-00		V	Nobivac Coryza S.C.0.5ml				
84	12	24-Mar-00		V	AE + POX + Dil: Wing Web - full				
112	16	21-Apr-00		V	Nobivac Coryza S.C.0.5ml				
112	16	22-Apr-00		V	Nobilis IB + ND: I.M.0.5ml				
119	17	29-Apr-00		V	Ma5+Clone30: Spray Vaccination				
119	17	8-May-00		V	BLOOD TESTING AND WATER ANALYSIS				

KEY
V

VACCINATION

RECCOMENDATION:

ALTERNATION BETWEEN VH+120 and MA5+CLONE30 Vaccines for NCD+IBV DISEASES

VH+120 VACCINATION FOR NCD+IBV AT : 30weeks, 42weeks, 66weeks

MA5+CLONE30 VACCINATION FOR NCD+IBV AT : 36weeks, 48weeks, 60weeks

DEWORM AT 10 AND 16 WEEKS

NOTE: THIS VACCINATION PROGRAM IS JUST A GUIDE AND IT IS SUSCEPTIBLE TO CHANGES ACCORDING TO THE FARMER'S POULTRY VETERINARIAN CONSULTANT

HYBRID POULTRY FARM DOES NOT ACCEPT ANY RESPONSIBILITY IN CASE OF ANY POULTRY DISEASE CHALLENGE OCCURRING